

# Strep-Tactin® Purification

## - Short Protocol -

### Purification of Strep-tag®II and Twin-Strep-tag® fusion proteins with Strep-Tactin® matrices

#### Strep-Tactin® affinity purification

The **Strep-tag®** purification system is based on the highly selective binding of engineered streptavidin, called Strep-Tactin®, to Strep-tag®II fusion proteins. This technology allows one-step purification of almost any recombinant protein under physiological conditions, thus preserving its biological activity. The Strep-tag® system can be used to purify functional Strep-tag®II proteins from any expression system including baculovirus, mammalian cells, yeast, and bacteria. Unique Strep-Tactin® affinity columns have been developed for this purpose and the corresponding operating protocol is described below. Strep-tag®:Strep-Tactin® affinity purification should not be performed discontinuously in batch mode which would result in significantly reduced protein purity and yield in comparison to column chromatography. Further, prolonged batch incubations increase the risk of proteolytic degradation of the target protein including cleavage of the tag. Because of its small size, Strep-tag® generally does not interfere with the biological activity of the fusion partner. Thus, removal of the tag becomes superfluous. Comprehensive reviews and scientific publications giving an overview of various Strep-tag® applications are listed at [www.strep-tag.com](http://www.strep-tag.com).

The **Twin-Strep-tag®** is a dimeric version of the Strep-tag®II and therefore binds with the same selectivity to Strep-Tactin® but with a higher affinity. This increased affinity allows the purification of Twin-Strep-tagged proteins even from batch or cell culture supernatants with excellent yields. In addition the Twin-Strep-tag® tolerates higher amounts of detergents and salts in buffers compared to Strep-tag®II. Since the overall conditions for Strep-tag®II and Twin-Strep-tag® are the same, the following protocol can be used for both tags.

#### Short Protocol of the Strep-Tactin® chromatography cycle

Perform all operations at a temperature amenable to the stability of your recombinant protein (between 4°C and 30°C). To achieve optimal purification results, comply with the specified volumes and their ratios (column bed, washing volumes etc., see page 3). At low expression levels, increase applied cell extract volumes to take advantage of the column capacity, without changing other volumes.


**Do not use this protocol for Strep-Tactin®XT matrices!**

#### For research use only

Strep-Tactin® is covered by US patent 6,103,493. Further patent applications are pending world-wide. Purchase of reagents related to these technologies from IBA provides a license for non-profit and in-house research use only. The use of this technology for commercial use requires a separate license from IBA. A license may be granted by IBA on a case-by-case basis, and is entirely at IBA's discretion. Please contact IBA for further information on licenses for commercial use. Strep-tag® and Strep-Tactin® are registered trademarks of IBA GmbH.


# Strep-Tactin® Purification

## - Short Protocol -



**Equilibrate the column**

**1. Add 2 CV (column bed volume) Buffer W**




**Apply the protein extract**

**2. Add 0.5 up to 10 CV cell extract**

(Frozen cell extracts have to be centrifuged after thawing (14.000 rpm, 5 min, 4 °C) in order to remove any aggregates that may have formed. Apply the cleared extract to the column)


**Wait until the cell extract has completely entered the column**



**Wash column**

**3. Wash column 5x with 1 CV of Buffer W**


Collect the wash fractions (1 CV each) and apply 2 µl of each subsequent wash fraction to an analytical SDS-gel (additionally apply 2 µl of the lysate and 2 µl of the flow through).



**Elute protein**

**4. Add 6x of 0.5 CV Buffer E**


Collect the eluate in 0.5 CV fractions. To get most of Strep-tag® fusion protein in one fraction, i.e. E2, only, add 0.8 CV (E1), then 1.4 CV (E2) and finally again 0.8 CV (E3). 20 µl samples of each fraction can be used for SDS-PAGE analysis.



**Regenerate column**

**5. Wash the column 3x with 5 CV Buffer R**


The color change from yellow to red indicates the regeneration process. The intensity of the red color is an indicator for the column activity status for the next round of purification.



**Remove HABA and equilibrate**

**6. Remove Buffer R by adding 2x 4 CV Buffer W (pH 8)**

This step is essentially the same for Strep-Tactin Sepharose®, Macroprep® or Superflow® (high capacity). If the column is not completely discolored after the removal step please wash with Buffer W at pH 10.5. Afterwards the columns have to be stored in **Buffer W at pH 8**.



**Store column**

**7. Store the column at 4 °C overlaid with 2 ml Buffer W (pH 8).**

### Recommended volumes for working with Strep-Tactin® columns

| Column bed volume (CV) | Protein extract volume* |                 | Wash buffer volume | Elution buffer volume |
|------------------------|-------------------------|-----------------|--------------------|-----------------------|
|                        | Strep-tag®II            | Twin-Strep-tag® |                    |                       |
| 0.2 ml                 | 0.1 – 2 ml              | 0.1 – 20ml      | 5 x 0.2 ml         | 6 x 0.1 ml            |
| 1 ml                   | 0.5 – 10 ml             | 0.5 – 100 ml    | 5 x 1 ml           | 6 x 0.5 ml            |
| 5 ml                   | 2.5 – 50 ml             | 2.5 – 500 ml    | 5 x 5 ml           | 6 x 2.5 ml            |
| 10 ml                  | 5 – 100 ml              | 5 – 1000 ml     | 5 x 10 ml          | 6 x 5 ml              |

**Table 1.** Recommended buffer volumes for chromatography on Strep-Tactin® columns

\*Adjust protein extract volume according to binding capacity of the column (please refer to the appropriate data sheet) and apply the extract as concentrated as possible in the recommended volume range. Note that these volumes are average values which can be different for certain proteins.

### Biotin in cell culture media

Please note that biotin binds with high affinity to Strep-Tactin® thereby efficiently competing binding of Strep-tag®II and Twin-Strep-tag®. This binding is in addition irreversible and does not allow regeneration of the Strep-Tactin® column (in contrast to bound desthiobiotin).

Especially media for mammalian cell or insect cell cultivation may contain significant amounts of biotin. Thus, if proteins are secreted to the culture medium, biotin must be masked by the addition of avidin (or biotin should be removed by dialysis or gel filtration) prior to Strep-Tactin® chromatography. For more information, particularly for a list enumerating the biotin content of different cell culture media, please refer to [www.iba-lifesciences.com](http://www.iba-lifesciences.com).

### Reagents compatible with the Strep-tag®:Strep-Tactin® interaction

For an overview of reagents, which have been successfully tested for the purification of Strep-tag®II and Twin-Strep-tag® fusion proteins and their respective concentrations, see [www.iba-lifesciences.com](http://www.iba-lifesciences.com).

For a more detailed protocol and troubleshooting please download the comprehensive Strep-Tactin® Purification Manual and other related protocols under [www.iba-lifesciences.com](http://www.iba-lifesciences.com).

**Do not use this protocol for Strep-Tactin®XT matrices!**



Australian distributors:  
Fisher Biotec Australia  
free call: 1800 066 077  
email: [info@fisherbiotec.com](mailto:info@fisherbiotec.com)  
web: [www.fisherbiotec.com](http://www.fisherbiotec.com)



Solutions  
For Life Sciences